

FIXED POINT SETS FOR PERMUTATION MODULES

PETER COLLINGS

Let $A = kG$, the group algebra of some finite group where the characteristic of the field k divides $|G|$. In contrast to working over the complex field, the kG -modules are not usually semisimple. If a Sylow p -subgroup of G is not cyclic then there are infinitely many indecomposable kG -modules, and we usually enjoy little control over the category of such modules. It is therefore an important problem to find classes of modules which may be expressed as a sum of a not very great number of indecomposables, and to understand the structure of these indecomposables.

Permutation kG -modules, and their indecomposable summands (called p -permutation modules), provide one such class. It is only necessary to consider transitive permutation actions and for a fixed group this class is a finite set. Furthermore each indecomposable summand M determines three invariants, namely a vertex, a quotient of a subgroup of G and a projective indecomposable p -permutation module for this quotient; and these three invariants characterise M up to isomorphism. These invariants were introduced by Green [3] and Brauer and adapted by Puig and, in the case of permutation modules, by Broué [1].

Given an indecomposable kG -module M , a vertex of M is a subgroup $H \leq G$ minimal subject to satisfying the condition that there be a kH -module L for which M is a component of $\text{Ind}_H^G(L)$. This vertex is unique up to conjugation in G . Any Sylow p -subgroup of G satisfies this condition and so vertices are always p -subgroups. The kH -module L is called a source of M and is unique up to conjugation in $N_G(H)$. However, M is a p -permutation module if and only if M is a component of a permutation module and this happens exactly when L is the trivial kH -module. Hence p -permutation modules are also known as trivial source modules (cf. [6, p216]).

The Brauer correspondent M^H of M is an indecomposable $k(N_G(H)/H)$ -module. When M is a component of a permutation module $k\Omega$ then M^H is a component of $k\text{Fix}_\Omega H$, hence is a p -permutation module. Broué's correspondence (1.1) asserts a 1 – 1 correspondence between the projective components of $k\text{Fix}_\Omega H$ and the components of M with vertex H .

In Theorem (2.11) we will seek to phrase Broué's correspondence in terms not of vertices but of *fixed point sets* (defined in (2.4)), a notion which we will see is equivalent to that of vertices.

In Section 3 and onwards we shall let n be a positive integer and utilize fixed point sets in the situation of permutation modules afforded by the action by conjugation of the symmetric group $\text{Sym}(qn)$ on its conjugacy class of fixed point free elements of order q . Our aim will be to determine as far as possible the fixed point sets of the components involved. At the same time we will show how the corresponding vertices may be computed eg Lemma (4.5) and Lemma (4.6), and (5.4).

In Theorem (7.18) we determine the general form of a fixed point set. From this theorem a list can be derived of possible fixed point sets in terms not just of irreducible sets but of the even more basic transitive sets. For a given p and q we need the further information of: (1) those values of n for which our permutation module admits a projective summand and (2) the numbers $\kappa(X)$ (defined in (7.15)) for irreducible fixed point sets X . These missing values are not investigated in the present paper.

When $p = q = 2$ the problem becomes to determine the fixed point sets, or the vertices of the components of, the permutation module afforded by the action of $\text{Sym}(2n)$ on its conjugacy class Ξ_{2n}^2 of fixed point free involutions. This permutation action has been studied before; as an example the ordinary character it affords is known, and we refer to [4] for a demonstration of this and associated results. In the situation $p = q = 2$ the values labelled (1) and (2) above can be obtained, leading to a precise list of vertices. In the author's PhD thesis [2] this precise list is constructed, along with an explicit determination of the number of components and the Brauer quotients. These results have been collected by the author elsewhere, along with a determination of the Brauer characters of the components, again using a result on ordinary characters contained in [4].

The tool to achieve the above is to view the natural embeddings of the cartesian product of any two symmetric groups into a third as a type of binary multiplication which will allow us to construct new fixed point sets from existing ones. The binary multiplication even allows for an essentially unique decomposition into irreducibles, with products of coprime fixed point sets being easy to control. Furthermore a unary multiplication is provided by the diagonal subset of the cartesian product.

1. PRELIMINARIES.

Let G be a finite group and let P be a Sylow p -subgroup of G . A kG -module has a P -invariant basis if and only if it is a component of some permutation kG -module, and such a module is called a p -permutation module. The main tool for identifying the vertices of the components of permutation modules is the following correspondence due to Broué [1, (3.2)]. The theorem following it (see [6, (27.7)]) is another useful characterisation of these vertices.

Broué Correspondence 1.1. *Let Ω be a permutation G -space. There is a multiplicity-preserving bijective correspondence between the components of the permutation kG -module $k\Omega$ with vertex $Q \leq G$ and the projective components of the permutation $k(N_G(Q)/Q)$ -module $k\text{Fix}_\Omega(Q)$.*

Theorem 1.2. *Let P be a Sylow p -subgroup of G . An indecomposable kG -module M with a P -invariant basis X has vertex conjugate to $Q \leq P$ if and only if: Q is a maximal subgroup of P subject to fixing an element of X .*

Recall that any transitive permutation kP -module is indecomposable (eg [6, (27.1)]). Thus the regular module kP is indecomposable and the only indecomposable projective kP -module is kP .

Lemma 1.3. *Let M be an indecomposable permutation P -module with a P -invariant basis X and let $x \in X$. Then $\text{Stab}_P(x)$ is a vertex of M . In particular P acts regularly on X if and only if M is projective.*

Proof. First of all we note that P must act transitively on X because M is indecomposable. A vertex of M is any subgroup Q of P maximal subject to fixing an element of X . By taking a conjugate of Q in P we may assume that Q fixes the element $x \in X$. Thus $Q \leq \text{Stab}_P(x)$ and the maximality of Q gives $Q = \text{Stab}_P(x)$. Now M is projective if and only if Q is trivial, and since $Q = \text{Stab}_P(x)$ this is the case if and only if P acts regularly on X . $\square$

The following is a standard result.

Lemma 1.4. Projective Modules. *Let M be a kG -module and P a Sylow p -subgroup of G . Then M is projective if and only if $\text{Res}_P^G(M)$ is projective.*

Proof. $\square$

Lemma 1.5. *A p -permutation kG -module M with P -invariant basis X is projective if and only if each P -orbit on X is regular.*

Proof. Let $X_1, X_2, \dots, X_s$ be the P -orbits on X . Then $\text{Res}_P^G(M) = kX_1 \oplus kX_2 \oplus \dots \oplus kX_s$ is a decomposition of $\text{Res}_P^G(M)$ into indecomposable summands. Now by Lemma (1.4) M is projective if and only if $\text{Res}_P^G(M)$ is projective, and by Lemma (1.3) this happens if and only if P acts regularly on each X_i . $\square$

The following is well known.

Proposition 1.6. *Let G and H be finite groups. Let $\{S_i\}_{i=1}^s$ be a complete set of pairwise non-isomorphic indecomposable projective kG -modules and $\{T_i\}_{i=1}^t$ a complete set of pairwise non-isomorphic indecomposable projective kH -modules. Then $\{S_i \otimes T_j\}_{1 \leq i \leq s, 1 \leq j \leq t}$ is a complete set of pairwise non-isomorphic indecomposable projective $k(G \times H)$ -modules.*

Corollary 1.7. *If G is a finite group and M is a kG -module let $np(M)$ be the number of indecomposable projective summands of M counting multiplicities when M is expressed as a direct sum of indecomposable modules. Then if G_1 and G_2 are finite groups and M_i is a kG_i -module for $i = 1, 2$ we have*

$$np(M_1 \otimes M_2) = np(M_1) \cdot np(M_2).$$

Proof. $\square$

Tensor Product with Wreath Product Action 1.8. Let M be a kG -module. Let a be a positive integer and let $G^a := G \times G \times \dots \times G$ be the direct product of a copies of G . Let $M^a := M \otimes M \otimes \dots \otimes M$ be the tensor product of a copies of M . Then M^a is a kG^a -module.

Now let $H := \text{Sym}(a)$. Then H can be made to act on G^a by permuting factors, or as it is sometimes called by *place permutations*. Thus for example we have $(g_1, g_2, \dots, g_a) \cdot (1\ 2) = (g_2, g_1, \dots, g_a)$. The action of H on G^a allows us to form the semidirect product $G^a \rtimes H$ which is known as the wreath product $G \wr H$. The subgroup $G^a = G \times G \times \dots \times G$ is called the base subgroup of G and the subgroup $H = \text{Sym}(a)$ is called the top subgroup of G .

Similarly we may define an action of H on the module M^a by letting H permute factors. This turns the tensor product M^a into a kH -module. Therefore M^a is

simultaneously a kG^a -module and a kH -module. We may define a $k(G \wr H)$ -module structure on M^a by stipulating

$$m^{gh} = (m^g)^h,$$

for $m \in M^a$, $g \in G^a$ and $h \in H$. We use M^{la} to denote this $k(G \wr H)$ module. In essence we let the subgroup G^a of $G \wr H$ act first via the product action and then we let the top subgroup $H \leq G \wr H$ act by permuting factors.

The above construction is performed for a kG -module M but is equally applicable to the situation of a permutation G -space X . If X is a permutation G -space then we define a permutation $G \wr H$ -space X^{la} as follows. As sets we let X^{la} be the cartesian product $X \times X \times \cdots \times X$ of a copies of X . Then we let G^a act via the product action and we let H act by permuting factors.

If $M = kX$ is the permutation kG -module afforded by X then $M^{la} = kX^{la}$ is the permutation module corresponding to the permutation $G \wr H$ -space X^{la} . In particular in this situation M^{la} is a permutation $k(G \wr H)$ -module.

The construction of the wreath product is of course standard and we refer to [5, (4.1)] for more details. Usually the wreath product $G \wr H$ of two groups G and H is defined in terms of a permutation G -space X and a permutation H -space Y . In the above construction we are taking $H = \text{Sym}(a)$ and the permutation H -space Y is the natural permutation $\text{Sym}(a)$ -space on the set of a letters $\{1, 2, \dots, a\}$.

Lemma 1.9. *Let M be a kG -module and let a be a positive integer. The $k(G \wr H)$ -module M^{la} admits a projective summand only if M admits a projective summand.*

Proof. We have $\text{Res}_{G^a}^{G \wr H}(M^{la}) = M^a$, so the kG^a -module M^a must admit a projective summand. The result now follows. $\square$

Lemma 1.10. *Let M be a kG -module and let $a < p$ be a positive integer. Suppose M admits a projective summand. Then the $k(G \wr H)$ -module M^{la} admits a projective summand.*

Proof. The size of $H = \text{Sym}(a)$ is coprime to p because $a < p$, so the base group G^a contains a Sylow p -subgroup of $G \wr H$. We know that the kG^a -module M^a admits a projective summand. Now $M^a = \text{Res}_{G^a}^{G \wr H}(M^{la})$ and we may apply (1.4) to infer that M^{la} admits a projective summand. $\square$

Lemma 1.11. *Let M be a kG -module. Assume $a \geq 2$ and M^{la} admits a projective summand as a kG^{la} -module. Then $M^{l(a-1)}$ admits a projective summand as a $kG^{l(a-1)}$ -module.*

Proof. For convenience let $M_i \cong M$ and $G_i \cong G$ for $i = 1, \dots, a$. Then $G^{la} \cong (G_1 \times \cdots \times G_{(a-1)} \times G_a) \cdot T$ where $T \cong \text{Sym}(a)$. So G contains the subgroup $G^{l(a-1)} \cong (G_1 \times \cdots \times G_{(a-1)} \times 1) \cdot T_1$ where $T_1 \leq T$ is the subgroup of T fixing the a -th factor of $M_1 \otimes \cdots \otimes M_a$ and $T_1 \cong \text{Sym}((a-1))$.

The restriction of M^{la} to this subgroup must also admit a projective summand and we have

$$\text{Res}_{G^{l(a-1)}}^{G^{la}}(M^{la}) \cong \bigoplus_{\dim | M |} M^{l(a-1)},$$

a direct sum of $\dim | M |$ isomorphic copies of $M^{l(a-1)}$.

It follows that the $kG^{(a-1)}$ -module $M^{(a-1)}$ must have a projective summand. $\square$

2. MAIN RESULTS ON FIXED POINT SETS.

In this section we will define the notion of *fixed point sets* for a permutation G -space Ω (over k); these will be in a bijective correspondence with the vertices of the components of the permutation kG -module $k\Omega$. The vertex corresponding to a given fixed point set is easily recovered, and vice versa, and we can reformulate Broué's correspondence in terms of this definition.

The maximality property of vertices requires that fixed point sets satisfy amongst general subsets of Ω a property which we will describe as *closed*. By restricting our attention to closed subsets of Ω we may replace in the Broué correspondence the quotient group $N_G(Q)/Q$ with another quotient group N_X/S_X which for our purposes is perhaps more convenient.

The next lemma describes how vertices - and thus projectivity - behave when we factor out p' -subgroups.

Lemma 2.1. *Let Ω be a permutation G -space and let $M := k\Omega$ be the corresponding permutation kG -module. Let N be a normal subgroup of G that fixes every element of Ω and let $\pi : G \rightarrow G/N$ be the quotient homomorphism. Let $M = M_1 \oplus M_2 \oplus \cdots \oplus M_t$ be a decomposition of M into indecomposable summands. Each summand M_i may be viewed as a kG -module or a $k(G/N)$ -module. A subgroup $Q \leq G$ is a vertex of a summand M_i as kG -modules if and only if $(Q)\pi$ is a vertex of M_i when viewed as a $k(G/N)$ -module.*

Proof. We may assume $Q \subset P$. Now P is a Sylow p -subgroup of G so $(P)\pi$ is a Sylow p -subgroup of G/N . Choose i and let X be a P -invariant basis of M_i . Then X is a $(P)\pi$ -invariant basis of M_i viewed as a $k(G/N)$ -module. If $Q \leq G$ is another subgroup of G then Q and $(Q)\pi$ fix the same elements of X , so the subgroup Q of P is maximal subject to fixing an element of X if and only if the subgroup $(Q)\pi$ of $(P)\pi$ is maximal subject to fixing an element of X . This concludes the proof. $\square$

Lemma 2.2. *Let G be a finite group and the finite set Ω be a transitive permutation G -space. Let Q be a vertex of some indecomposable summand of $K\Omega$ and let P be a Sylow p -subgroup of G that contains Q . Then there is some element $\omega \in \Omega$ such that $Q = P \cap \text{Stab}_G(\omega)$.*

Proof. Let $k\Omega = M \oplus M'$ where M is indecomposable with vertex $Q \leq P$. We have $\text{Res}_P^G(k\Omega) = \bigoplus_{i=1}^t k\Omega_i$ where the Ω_i are the P -orbits in Ω . Note that the $k\Omega_i$ are indecomposable as kP -modules since P is a p -group. By Krull-Schmidt we have therefore that

$$\text{Res}_P^G(M) = \bigoplus_{i \in I}^t k\Omega_i$$

where I is some subset of $\{1, 2, \dots, t\}$. Identify M as a P -module with $\bigoplus_{i \in I}^t k\Omega_i$. Then M has P -invariant basis $X = \bigcup_{i \in I} \Omega_i$. Let $Q \leq P$ be a vertex of M . Then

by Theorem 1.2, Q is a maximal subgroup of P subject to fixing an element of X . So there is some $\omega \in \Omega_i$ for some i which is fixed by Q but not by any subgroup R of P properly containing Q .

Then $P \cap \text{Stab}_G(\omega) = \text{Stab}_P(\omega) = Q$. $\square$

Theorem 2.3. *Let M be a permutation kG -module, say $M = k\Omega$. Let Q be a vertex of an indecomposable summand of M , let X be the set of fixed points of Q on Ω , and let S be the pointwise stabilizer of X in G . Then Q is a Sylow p -subgroup of S .*

Proof. Let $R \geq Q$ be a Sylow p -subgroup of S . If $R \neq Q$ let $U := N_R(Q)$. Then U is a p -subgroup which strictly contains Q and we have

$$Q < U \leq N_G(Q) \cap S.$$

The factor group $N_G(Q)/Q$ therefore contains the non-trivial p -subgroup $\overline{U} := U/Q$. But $U \leq S$ so U and $\overline{U}$ act trivially on F . Certainly then a Sylow p -subgroup of $N_G(Q)/Q$ cannot act regularly on any element of F , and by Lemma (1.3) the $kN_G(Q)/Q$ -module kF does not admit a projective summand. Of course by the Broué correspondence (1.1) this implies that Q is not a vertex of an indecomposable summand of M . The alternative $Q = R$ must be the case. $\square$

Notation 2.4. Suppose Ω is a G -set. Then if $X \subset \Omega$ is a subset we define $S_X := \text{Stab}_G(X)$ to be the pointwise stabilizer, and we write Q_X for a Sylow p -subgroup of S_X . We will also write N_X for the set stabilizer of X in G , that is, $N_X := \{g \in G \mid X^g = X\}$.

Definition 2.5. Suppose Ω is a G -set and $X \subseteq \Omega$. We say that X is a fixed point set if there is a vertex of some indecomposable summand of $k\Omega$ with vertex Q such that $X = \text{Fix}_\Omega(Q)$.

Theorem 2.3 shows that in this case Q is a Sylow p -subgroup of S_X and so for a fixed point set we may take $Q_X = Q$.

We can reformulate part of the Broué correspondence as follows:

Corollary 2.6. *The set $X \subseteq \Omega$ is a fixed point set if and only if the p -permutation module kX for $N_G(Q_X)/Q_X$ has a projective summand.*

Proof. $\square$

Definition 2.7. Suppose Ω is a G -set, and $X \subseteq \Omega$ is a subset. We say that X is *closed* if $\text{Fix}_\Omega(Q_X) = X$. This notion is independent of the choice of Q_X . Note that if X is a fixed point set, then X is closed.

One way of obtaining closed subsets is to take the closure of a given subset. For subgroups L of G we denote by $*$ an assignment (depending on a choice of P) $L \mapsto \text{Fix}_\Omega(\text{Syl}_p(L))$ then we mean that L^* is the set of fixed points in Ω of some Sylow p -subgroup of L . The reverse assignment is to denote by $*$ a choice of map $X \mapsto \text{Syl}_p(\text{Stab}_G(X))$ which assigns to the subset X of Ω some Sylow p -subgroup X^* of the stabilizer in G of X . We may say that the subset $X^{**} \subset \Omega$ is the *closure* of X . Thus X is closed exactly when $X = X^{**}$, and this must happen when for instance X is a fixed point set. Similarly for a vertex Q of some component of $k\Omega$ we must have $Q = Q^{**}$.

Properties of Closed Subsets 2.8. Suppose X is closed, that is $\text{Fix}_\Omega(Q_X) = X$. Then we even have $\text{Fix}_\Omega(S_X) = X$. In general, the normalizer of a subgroup

permutes the fixed points of this subgroup, so the normalizer $N_G(S_X)$ also leaves X invariant. We have defined (in (2.4))

$$N_X := \{g \in G : (X)g \subseteq X\}$$

the largest subgroup of G that leaves X invariant. In general, $N_X \subseteq N_G(S_X)$. When X is closed, we therefore have $N_G(S_X) = N_X$. Our aim now is to characterize fixed point sets in terms of N_X and S_X (which will allow us to characterize vertices in terms of fixed point sets).

Notation 2.9. Let Ω be a G -set and let $X \subset \Omega$ be a closed subset. Define

$$M_X := N_X/S_X.$$

As we mentioned above, the group N_X acts on X with kernel S_X . Thus M_X acts on X and X is a faithful permutation M_X -space. The action of N_X on X induces a homomorphism $N_X \rightarrow \text{Sym}(X)$, and M_X may be taken to be the image of this homomorphism.

Proposition 2.10. *Given a closed subset $X \subset \Omega$ we have $N_G(Q_X) \leq N_X$, and also $N_G(Q_X) \cdot S_X = N_X$.*

Proof. The subgroup $N_G(Q_X)$ will leave invariant the set $\text{Fix}_\Omega(Q_X)$; but X is closed so this set is again X , whence $N_G(Q_X)$ leaves X invariant giving $N_G(Q_X) \leq N_X$.

That $N_G(Q_X) \cdot S_X = N_X$ follows from a Frattini argument since Q is a Sylow p -subgroup of S_X and $N_G(Q_X) \leq N_X = N_G(S_X)$. (The argument attributed to Frattini is that if P is a Sylow p -subgroup of a normal subgroup N of G then $G = N_G(P) \cdot N$.)

Theorem 2.11. *Let $X \subset \Omega$ be closed. Then the projective components of the kM_X -module kX are in bijective correspondence with the projective components of the $k(N_G(Q_X)/Q_X)$ -module kX .*

Proof. By the preceding proposition we have

$$\begin{aligned} M_X &= N_X/S_X = N_G(Q_X) \cdot S_X/S_X \\ &\cong N_G(Q_X)/N_{S_X}(Q_X) \cong (N_G(Q_X)/Q_X)/(N_{S_X}(Q_X)/Q_X). \end{aligned}$$

Now $N_{S_X}(Q_X)/Q_X$ has p' size since Q_X is a Sylow p -subgroup of S_X . The result follows from Lemma (2.1). $\square$

We can now reformulate the Broué Correspondence:

Theorem. 2.12. **Broué's correspondence for closed sets** Let $X \subset \Omega$ be closed. There is a multiplicity-preserving bijective correspondence between the isomorphism types of components of $k\Omega$ which admit X as a fixed point set and the set of projective components of the permutation kM_X -module kX (up to isomorphism).

Corollary. 2.13. *Let $X \subset \Omega$ be closed. Then X is a fixed point set if and only if the permutation kM_X -module kX admits a projective component.*

3. SYMMETRIC GROUPS AND CONJUGACY ACTIONS.

A Specific Problem. We have seen that the problem of determining the vertices of the indecomposable summands of permutation modules of finite groups is equivalent to finding the corresponding fixed point sets. Now we shall study these fixed point sets for the permutation $k\text{Sym}(qm)$ -module $k\Xi_{qm}^q$ where Ξ_{qm}^q is the conjugacy class of $\text{Sym}(qm)$ consisting of fixed point free elements which are products of q -cycles and k is a field of characteristic p .

Irreducible Sets 3.1. In such actions, that is when a group G acts on itself by conjugation, the subsets of the relevant G -set have an additional structure given by the group multiplication. We will therefore consider ways of building new fixed point sets out of known ones, using the provided multiplication, and also using the fact that the direct product of two symmetric groups can be embedded into a larger symmetric group.

In the first part of this section we will study set constructions, and at the end of the section we will relate these set constructions to fixed point sets.

Definition 3.2. The group of finitary permutations of $\mathbb{N}$ is defined to be the union of all finite symmetric groups:

$$\text{Sym}_f(\mathbb{N}) := \bigcup_{n=1}^{\infty} \text{Sym}(n).$$

Recall that the support of a permutation ρ is defined as:

$$\text{supp } \rho := \{\omega \in \mathbb{N} \mid \omega(\rho) \neq \omega\}.$$

If X is a set of finitary permutations we set $\text{supp}(X) = \bigcup_{\rho \in X} \text{supp}(\rho)$. Now we define $\mathcal{F}$ to be all finite subsets of $\text{Sym}_f(\mathbb{N})$, excluding the empty set and the set which contains only the identity permutation. That is,

$$\mathcal{F} := \{X \subset \text{Sym}_f(\mathbb{N}) : 0 < |X| < \infty \text{ and } X \neq \{1\}\}.$$

For an integer $q \geq 2$, we define $\mathcal{S}^q \subset \mathcal{F}$ to be all subsets X such that

- (1) each $\rho \in X$ is a product of q -cycles;
- (2) for each $\rho \in X$ we have $\text{supp}(\rho) = \text{supp}(X)$.

Any such $X \in \mathcal{S}^q$ is conjugate to a subset of Ξ_{qm}^q where qm is the size of $\text{supp}(X)$.

Notation 3.3. Let X be an element of $\mathcal{F}$. By construction X is a subset of some finite subgroup of $\text{Sym}(\mathbb{N})$, and in fact we have $X \subset \text{Sym}(\text{supp } X) \leq \text{Sym}(\mathbb{N})$. So let $G_X := \text{Sym}(\text{supp } X)$, and write $N_X := N_{G_X}(X) = \{g \in G_X \mid X^g = X\}$ and $S_X := \text{Stab}_{G_X}(X)$.

Now S_X is a normal subgroup of N_X and N_X acts by conjugation on X with kernel S_X . Thus $M_X := N_X/S_X$ acts on X with trivial kernel and kX is a faithful permutation kM_X -module. Let $Q_X \leq S_X$ be a Sylow p -subgroup of S_X .

These definitions were also made in Section (2.4), and it was suggested there that a distinctive case is when the permutation kM_X -module kX admits a projective summand. This is the condition which characterizes, amongst closed sets, the fixed point sets.

Multiplication In $\mathcal{F}$ 3.4. Let $X, Y \in \mathcal{F}$. Letting $a := |\text{supp } X|$ and $b :=$

$|\text{supp } Y|$, there is an embedding of groups

$$\tau : \text{Sym}(a) \times \text{Sym}(b) \rightarrow \text{Sym}(a + b).$$

We identify $\text{Sym}(a)$ with $\text{Sym}(\text{supp } X)$ and $\text{Sym}(b)$ with $\text{Sym}(\text{supp } Y)$ so that this embedding restricts to an embedding of the Cartesian product of X and Y :

$$\tau : X \times Y \rightarrow \text{Sym}(a + b).$$

We write $X * Y$ to mean the image of this embedding. By convention we write $X * \{1\} = X$ for any subset $X \in \mathcal{F}$. Now consider the s -fold Cartesian product $X^s = X \times X \times \cdots \times X$ and the embedding $\tau : X^s \rightarrow \text{Sym}(as)$. We let $\Delta^s X$ be the image under τ of the diagonal subset $\{(x, x, \dots, x) \mid x \in X\}$ of $(X)^s$.

We write the **degree** of Y as $d(Y)$ and define it by $d(Y) := |\text{supp } Y|$.

Remark 3.5. We will say that two elements X and Y of $\mathcal{F}$ are equivalent if there is $\sigma \in \text{Sym}(\mathbb{N})$ such that $X^\sigma = Y$. The $*$ -product can be thought of as defined on the resulting equivalence classes and in this case the $*$ -product is abelian and associative.

Definition 3.6. Irreducible Sets. We say that $X \in \mathcal{F}$ is **irreducible** if there do not exist elements $Y, Z \in \mathcal{F}$ such that $X = Y * Z$. Clearly X can be written as a product of irreducible members of $\mathcal{F}$. Note that $X \in \mathcal{S}^q$ is irreducible if and only if there do not exist in $Y, Z \in \mathcal{S}^q$ such that $X = Y * Z$.

Remark 3.7. If we interpret the product $*$ as being defined on the set of equivalence classes of $\mathcal{F}$ then Lemma 3.9 immediately implies that any decomposition into irreducible factors is unique. In fact slightly more is implied because the statement " $Y_i = Z_i$ " which forms part of the statement of the lemma asserts an absolute equality, that is, as subsets of $\text{Sym}(\mathbb{N})$, and not just equivalence.

Lemma 3.8. *The set $Y \in \mathcal{S}^q$ is a fixed point set if and only if each of the following hold*

- (1) Y is closed
- (2) the permutation kM_Y -module kY admits a projective summand.

Proof. This is a combination of Corollary (2.13) and Theorem (2.3). $\square$

Lemma 3.9. Uniqueness of Irreducible Decomposition. *Let $X \in \mathcal{F}$ be such that $X = Y_1 \times Y_2 \times \cdots \times Y_s = Z_1 \times Z_2 \times \cdots \times Z_t$ where the Y_i and Z_i are irreducible elements of $\mathcal{F}$. Then $s = t$ and after a reordering of factors we have $Y_i = Z_i$ for $1 \leq i \leq s$.*

Proof. We may assume that respectively the Y_i and the Z_i have pairwise disjoint supports and that $X = Y_1 \times Y_2 \times \cdots \times Y_s = Z_1 \times Z_2 \times \cdots \times Z_t$. Put $s_i := \text{supp } Y_i$ and $t_i := \text{supp } Z_i$. Now X is a subset of the subgroup $\text{Sym}(s_1) \times \text{Sym}(s_2) \times \cdots \times \text{Sym}(s_s) \leq \text{Sym}(\mathbb{N})$ and we may consider the projection $\pi_{Y_i} : X \rightarrow Y_i \subset \text{Sym}(s_i)$, and similarly the projection $\pi_{Z_i} : X \rightarrow Z_i$. Thus we have $Y_1 = (X)\pi_{Y_1} = (Z_1 \times Z_2 \times \cdots \times Z_t)\pi_{Y_1} = (Z_1)\pi_{Y_1} \times (Z_2)\pi_{Y_1} \times \cdots \times (Z_t)\pi_{Y_1}$ and since Y_i is irreducible this forces $Y_1 = (Z_i)\pi_{Y_1}$ for some i , and $s_1 \subset t_i$.

Similarly we have $Z_i = (Y_1)\pi_{Z_i} \times (Y_2)\pi_{Z_i} \times \cdots \times (Y_s)\pi_{Z_i}$. But $s_1 \subset t_i$ so $(Y_1)\pi_{Z_i} = Y_1$, and the irreducibility of Z_i implies $Z_i = Y_1$. Reordering the Z_j we

have $Z_1 = Y_1$; upon cancelling these two factors the result follows by induction. $\square$

Let $N_Y = \{g \in \text{Sym}(\text{supp } Y) : Y^g = Y\}$ for $Y \in \mathcal{F}$.

Corollary 3.10. *Let $Y \in \mathcal{F}$ and let $Y = Y_1 \times Y_2 \times \cdots \times Y_s$ be a decomposition into irreducibles. For every $1 \leq i \leq s$ and $g \in N_Y$ there is $1 \leq j \leq s$ such that $Y_i^g = Y_j$.*

Proof. We have $Y = Y^g$ and $Y_1 \times Y_2 \times \cdots \times Y_s = Y_1^g \times Y_2^g \times \cdots \times Y_s^g$ so the result follows immediately from the lemma. $\square$

Corollary 3.11. *Then the action of N_Y on Y induces a permutation N_Y -action on the set of irreducible factors of Y .*

Proof. This is a restatement of the previous corollary. $\square$

Lemma 3.12. *Suppose that $X \in \mathcal{S}^q$ is irreducible and satisfies $|X| > 1$. Then $\Delta^i X$ is irreducible for every i .*

Proof. We write $\Delta^i X = X_1 \Delta X_2 \Delta \cdots \Delta X_i$ where each X_j is conjugate to X and where the $\text{supp } X_j$ are pairwise disjoint. Put $\alpha := \bigcup_{j=1}^i \text{supp } X_j$. Assume that $\Delta^i X = Y \times Z$ where Y and Z are elements of $\mathcal{S}^q$ with disjoint non-empty support. Then for $1 \leq j \leq i$ we have $X_j = (\Delta^i X)\pi_{X_j} = (Y)\pi_{X_j} \times (Z)\pi_{X_j}$, and since X_j is irreducible this implies that either $X_j = (Y)\pi_{X_j}$ or $X_j = (Z)\pi_{X_j}$. Consequently either $\text{supp } X_j \subset \text{supp } Y$ or $\text{supp } X_j \subset \text{supp } Z$.

Let a be the set of j such that $\text{supp } X_j \subset \text{supp } Y$ and let b be the set of j such that $\text{supp } X_j \subset \text{supp } Z$. Since $\text{supp } Y$ and $\text{supp } Z$ are non-empty it follows that a and b are each non-empty. Also, since $\alpha = \text{supp } Y \cup \text{supp } Z$ we have $\text{supp } Y = \bigcup_{j \in a} \text{supp } X_j$ and $\text{supp } Z = \bigcup_{j \in b} \text{supp } X_j$. After a possible reordering we may assume that $a = \{1, 2, \dots, s\}$ and $b = \{s+1, s+2, \dots, i\}$.

Comparing this with $X_1 \Delta X_2 \Delta \cdots \Delta X_i = Y \times Z$ we have $Y = (Y \times Z)\pi_Y = X_1 \Delta X_2 \Delta \cdots \Delta X_s$ and similarly $Z = X_{s+1} \Delta X_{s+2} \Delta \cdots \Delta X_i$. Thus $|Y| = |Z| = |X|$ and $|Y \times Z| = |X|^2$. But $|Y \times Z| = |\Delta^i X| = |X|$, and so $|X| = 1$. $\square$

Conjugation actions. 3.13. Let $X \in \mathcal{F}$, and define

$$G_X := \text{Sym}(\text{supp } X),$$

the symmetric group permuting the elements in $\text{supp } (X)$.

We view X as a subset of a permutation G -set, where G acts by conjugation. That is,

$$\Omega = \Xi_X := \bigcup_{g \in G} X^g \subseteq G_X.$$

We observe that in this setup S_X is equal to $C_{G_X}(X)$ and N_X is equal to $N_{G_X}(X)$. The set of fixed points of Q_X on Ω is then equal to

$$\text{Fix}_\Omega(Q_X) = C_{G_X}(X) \cap \Omega.$$

Note that we always take these with respect to the G_X -set just defined.

Conventions 3.14. Now we specialize these concepts to $Y \in \mathcal{S}^q$. Recall then that Y is a fixed point set if there is an indecomposable summand of the permutation kG_Y -module $k\Xi_Y$ with vertex $Q \leq G_Y$ such that $Y = \text{Fix}_{\Xi_Y}(Q)$. Thus Y is a fixed point set if and only if Q_Y is a vertex of an indecomposable summand of the permutation module $k\Xi_Y$, for the group G_Y . For example, if

$Y \in \mathcal{S}^q$ is such that all elements of Y have the same support then Ξ_Y is actually the G_Y set of all fixed point free elements which are the product of q -cycles.

Exact Sets and Projective Sets 3.15. Let $Y \in \mathcal{S}^q$. We say that Y is **exact** if Q_Y is fixed point free in its action on $\text{supp } Y$, that is, if $\text{supp } Q_Y = \text{supp } Y$. Recall that for $Y, Z \in \mathcal{S}^q$ we have made choices so that $Q_X * Q_Y \leq Q_{X*Y}$, and it follows that the product of exact sets is exact.

On the other hand, we say that Y is **projective** if $Q_Y = 1$. We observe that $Y \in \mathcal{S}^q$ is projective and closed if and only if $Y = \Xi_Y$.

Lemma 3.16. *Suppose that $Y \in \mathcal{S}^q$ is irreducible and closed. Then X is exact or X is projective.*

Proof. Let $\alpha = \text{supp } Q_X$ and write $\text{supp}(X) = \alpha \cup \beta$ where Q_X fixes each element of β . The group $\langle X \rangle$ centralizes Q_X and therefore it permutes the orbits of Q_X on the support of X . Assume for a contradiction that α and β are both $\neq \emptyset$. Then $X \subset \text{Sym}(\alpha) \times \text{Sym}(\beta)$.

So we have $X \subseteq X_\alpha * X_\beta$ (where X_α is the set of all factors of X which are supported on α).

By assumption, X is closed, that is $X = \text{Fix}_{\Omega_X}(Q_X)$.

We can describe these fixed points: they are all elements in Ω_X which lie in $\text{Sym}(\alpha) \times \text{Sym}(\beta)$, and where the factor in $\text{Sym}(\alpha)$ commutes with Q_X . The factor in $\text{Sym}(\beta)$ is arbitrary. In fact we get $X = Y * X_\beta$ where

$$Y = \text{Fix}_{\Omega_{X_\alpha}}(Q_X), X_\beta = \Omega_{X_\beta}.$$

We assume X is irreducible, so this is a contradiction. $\square$

Remark. We have seen at the same time that if $Q_X = 1$ then $X = X_\beta$ and this is the complete G_X -set Ω_X . This will be relevant later.

4. PERMUTATION STRUCTURE OF PRODUCTS OF IRREDUCIBLE SETS.

Suppose that we have found a fixed point set $Y \in \mathcal{S}^q$ which is irreducible and exact. By Lemma (3.8) Y is closed and the permutation kM_Y -module kY admits a projective summand. We wish to find conditions under which the elements $*^s Y$ and $\Delta^s Y$ are fixed point sets. This means that (i) we want find conditions under which $*^s Y$ or $\Delta^s Y$ is a closed element of $\mathcal{S}^q$, and (ii) we would like to know the structure of the permutation $kM_{*^s Y}$ -module $k*^s Y$ and of the permutation $kM_{\Delta^s Y}$ -module $k\Delta^s Y$. We address the second of these first. In doing so we shall make use of the wreath product constructions described in Section (1.8).

Actions on Products 4.1. It is well known that the stabilizer in $\text{Sym}(qn)$ of an element of Ξ_{qn}^q (that is, a fixed point free element which is a product of q -cycles) is isomorphic to the wreath product $\mathbb{Z}_q \wr \text{Sym}(n)$. We now consider the version of this result which applies to the situation where we are dealing not with q -cycles in $\text{Sym}(qn)$ but equal sized subsets of $\{1, 2, \dots, qn\}$, or of some similar set.

Let α be a finite subset of $\mathbb{N}$ and let the subsets $\alpha_1, \alpha_2, \dots, \alpha_s$ partition α and satisfy $|\alpha_i| = |\alpha_1|$. Let $G := \text{Sym}(\alpha_1)$. Then α_1 is a natural permutation G -space and as described in Section (1.8) we may form the tensor product α_1^{ls} giving

a permutation $(G \wr \text{Sym}(s))$ -space. In fact this construction can be made inside $\text{Sym}(\alpha)$. The action of $\text{Sym}(\alpha)$ on α induces an action on sets of subsets of α and we may define C to be the stabilizer under this action of the set $A := \{\alpha_1, \alpha_2, \dots, \alpha_s\}$. It is reasonably clear that C is isomorphic to the wreath product $\text{Sym}(\alpha_1) \wr \text{Sym}(s)$. However, we shall now present the details of the proof.

Wreath Products 4.2. The base group in the situation above is the Young subgroup $B := \text{Sym}(\alpha_1) \times \text{Sym}(\alpha_2) \times \dots \times \text{Sym}(\alpha_s)$ and there is a canonical choice of subgroup $T \leq C, T \cong \text{Sym}(s)$ whose action on the set A is the full Symmetric action on s elements. This canonical choice is determined by the following. For each i, j there is a unique order preserving involution $g_{ij} \in \text{Sym}(\mathbb{N})$ such that $\text{supp } g_{i,j} = \alpha_i \cup \alpha_j$ and $\alpha_i^{g_{ij}} = \alpha_j$. We have $g_{ij} = g_{ji}$ and g_{ij} can be viewed as the transposition $(\alpha_i \alpha_j)$ in $\text{Sym}(A)$. Thus the g_{ij} generate a subgroup $T \cong \text{Sym}(A) \cong \text{Sym}(s)$ of C .

Note that although T is transitive on A , each of the above generators of T is order-preserving so the set of least elements of the α_i is left invariant by T , and T is not transitive on $\alpha = \alpha_1 \cup \dots \cup \alpha_s$ if $|\alpha_1| > 1$.

The subgroup $B \cdot T = \text{Sym}(\alpha_1) \cdot T \leq C$ is isomorphic to the wreath product $\text{Sym}(\alpha_1) \wr T$. If $g \in \text{Sym}(\alpha)$ is any other element such that $A^g = A$ then g induces some permutation on the elements of A and there is an element $t \in T = \text{Sym}(A)$ which is such that gt leaves each α_i invariant. Thus gt lies in the base group B and we have $C = B \cdot T \cong \text{Sym}(\alpha_1) \wr \text{Sym}(s)$.

Wreath Products and Irreducibles 4.3. If Y is an element of $\mathcal{S}^q$ then the product $*^s Y$ is equivalent to any product $Y_1 \times Y_2 \times \dots \times Y_s$ where each Y_i is conjugate in $\text{Sym}(\mathbb{N})$ to Y and where the sets $\text{supp } Y_i$ partition $\text{supp } *^s Y$. Thus by taking $\alpha_i := \text{supp } Y_i$ and $\alpha := \text{supp } *^s Y$ we are in the situation of the preceding paragraphs and there is a canonical choice of subgroup $T \cong \text{Sym}(s)$ in $G_{*^s Y} = \text{Sym}(\text{supp } *^s Y)$ which permutes the α_i . We may replace $Y_i \subset \text{Sym}(\alpha_i)$ with any subset of $\text{Sym}(\alpha_i)$ equivalent to Y_i . We may therefore assume that $Y_i = Y_1^{g_{1i}}$. The subgroup T now permutes the factors Y_i . In the following lemmas we shall assume that the subgroups T, B and C are as defined above (recall also the conventions given in (3.14) and (3.15)).

Lemma 4.4. *Let $Y \in \mathcal{S}^q$ be irreducible. Then*

$$N_{*^s Y} = (*^s N_Y) \cdot T \cong N_Y \wr \text{Sym}(s).$$

Proof. Write $*^s Y = Y_1 \times Y_2 \times \dots \times Y_s$ where each Y_i is conjugate in $\text{Sym}(\mathbb{N})$ to Y and where the $\text{supp } Y_i$ are pairwise disjoint. On one hand we clearly have $T \leq N := N_{*^s Y}$, and also $*^s N_Y \subseteq N$.

Now let $g \in N$. The irreducibility condition implies that g must permute the factors Y_i and so g must permute the subsets $\text{supp } Y_i$. In other words g must lie in the subgroup C . There is an element $t \in T \leq N$ such that gt leaves each $\text{supp } Y_i$ invariant. Thus $gt \in B \cap N = *^s N_Y$ and $N \leq (*^s N_Y) \cdot T$. Since $N_Y \leq N$ we have $N = (*^s N_Y) \cdot T \cong N_Y \wr \text{Sym}(s)$. $\square$

Lemma 4.5. *Let $Y \in \mathcal{S}^q$ be irreducible. Then*

$$S_{\Delta^s Y} = (*^s S_Y) \cdot T \cong S_Y \wr \text{Sym}(s).$$

Proof. Write $S := S_{\Delta^s Y}$. Then $T \leq S$ and $B \cap S = *^s S_Y$ so the proof of the last lemma works here. $\square$

If $Y \in \mathcal{S}^q$ contains exactly one element then $*^s Y = \Delta^s Y$. The next lemma assumes that this is not the case.

Lemma 4.6. *Let $Y \in \mathcal{S}^q$ be irreducible with $|Y| > 1$. Then*

$$S_{*^s Y} = *^s S_Y.$$

Proof. We may assume $s \geq 2$. Write as usual $*^s Y = Y_1 \times Y_2 \times \cdots \times Y_s$ and put $S := S_{*^s Y}$. Clearly we have $*^s S_Y \leq S$. Let a and b be distinct elements in Y and let $g \in S$. Then $\Delta^s Y \subset *^s Y$ so g certainly fixes $\Delta^s Y$, and $g \in S_{\Delta^s Y} = (*^s S_Y) \cdot T$, using the previous lemma. Thus $g = ut$ where $u \in *^s S_Y \leq S$ and $t \in T$, forcing $t \in S$. Suppose that $Y_1^t = Y_i$. If $i \neq 1$ then the element $a * b * \cdots * b \in *^s Y$, corresponding to the ‘coordinate vector’ with an a in the first place and b ’s elsewhere, is not centralized by t . It follows that $Y_1^t = Y_1$; similarly $Y_i^t = Y_i$ for each i and $t = 1$. Therefore $S \leq *^s S_Y$ as required. $\square$

Lemma 4.7. *Let $Y \in \mathcal{S}^q$ be irreducible and let $S := S_{\Delta^s Y}$. Then*

$$N_{\Delta^s Y} = \langle \Delta^s N_Y, S \rangle.$$

Proof. Put $N := N_{\Delta^s Y}$. Clearly $\Delta^s N_Y \leq N \leq C$ and $T \leq S \leq N$. Let $g \in N$; then $g = ut$ where $u \in B$ and $t \in T$. Since $t \in N$ we get $u \in N$. Writing all elements in $B = *^s G_Y$ as vectors with coordinates in G_Y we have $u = (u_1, u_2, \dots, u_s)$. If $y \in Y$ then $(y, y, \dots, y) \in \Delta^s Y$ and $(y, y, \dots, y)^u = (y^{u_1}, y^{u_2}, \dots, y^{u_s}) \in \Delta^s Y$. Therefore $y^{u_1} \in Y$ so $u_1 \in N_Y$. Also $y^{u_i} = y^{u_1}$ holds for every $y \in Y$ so $u_1 u_i^{-1} \in S_Y$ and $u = (u_1 u_1^{-1}, u_2 u_1^{-1}, \dots, u_s u_1^{-1})(u_1, u_1, \dots, u_1)$ gives $u \in \Delta^s N_Y \cdot *^s S_Y \leq \Delta^s N_Y \cdot S$, as needed. $\square$

Remark 4.8. As a corollary to this lemma we have $N_{\Delta^s Y}/S_{\Delta^s Y} \cong \Delta^s N_Y/(\Delta^s N_Y \cap S) = \Delta^s N_Y/\Delta^s S_Y \cong N_Y/S_Y \cong M_Y$. Thus $M_{\Delta^s Y} \cong M_Y$. We want to show slightly more than this, namely that the pair $(M_{\Delta^s Y}, \Delta^s Y)$ is permutation isomorphic to the pair (M_Y, Y) .

If $Y \in \mathcal{S}^q$ we shall let $\tau : G_Y \rightarrow \Delta^s G_Y$ be the bijection given by $g \mapsto (g, g, \dots, g)$.

Lemma 4.9. *Let $Y \in \mathcal{S}^q$ be irreducible. Then the pair $(M_{\Delta^s Y}, \Delta^s Y)$ is permutation isomorphic to the pair (M_Y, Y) .*

Proof. Let $\pi_\Delta : N_{\Delta^s Y} \rightarrow \text{Sym}(\Delta^s Y)$ be the homomorphism induced by the action of $N_{\Delta^s Y}$ on $\Delta^s Y$ and let $\pi_Y : N_Y \rightarrow \text{Sym}(Y)$ be the homomorphism induced by the action of N_Y on Y . By the previous lemma we may let $\iota : \Delta^s N_Y \rightarrow N_{\Delta^s Y}$ be the inclusion homomorphism. Then there is a homomorphism

$$\tau \iota \pi_\Delta : N_Y \rightarrow \Delta^s N_Y \rightarrow N_{\Delta^s Y} \rightarrow (N_{\Delta^s Y}) \pi_\Delta = M_{\Delta^s Y} \leq \text{Sym}(\Delta^s Y),$$

The homomorphism $\tau \iota \pi_\Delta : N_Y \rightarrow M_{\Delta^s Y}$ is surjective since by the previous lemma the kernel $S_{\Delta^s Y}$ of π_Δ together with $\Delta^s N_Y$ generates $N_{\Delta^s Y}$.

Now τ restricts to a bijection $\tau : Y \rightarrow \Delta^s Y$ which induces an isomorphism $\tau : \text{Sym}(Y) \rightarrow \text{Sym}(\Delta^s Y)$ and there is a homomorphism

$$\pi_Y \tau : N_Y \rightarrow \text{Sym}(Y) \rightarrow \text{Sym}(\Delta^s Y).$$

In fact for all $g \in N_Y$ the elements $(g) \tau \iota \pi_\Delta$ and $(g) \pi_Y \tau$ are each equal to that element of $\text{Sym}(\Delta^s Y)$ which takes $(y, y, \dots, y) \in \Delta^s Y$ to $(y^g, y^g, \dots, y^g) \in \Delta^s Y$. Thus

$\tau\iota\pi = \pi_Y\tau$. But $M_{\Delta^s Y} = N_{\Delta^s Y}\pi_\Delta = N_Y\tau\iota\pi_\Delta$ and $M_Y = N_Y\pi_Y$ so $M_Y\tau = M_{\Delta^s Y}$. $\square$

Lemma 4.10. *Let $Y \in \mathcal{S}^q$ be irreducible with $|Y| > 1$. Then the pair $(M_{*^s Y}, *^s Y)$ is permutation isomorphic to the pair $(M_Y \wr \text{Sym}(s), Y^{ls})$.*

Proof. Recall that the subgroup T satisfies $T \cong \text{Sym}(s)$. By Lemma (4.4) we have $N_{*^s Y} = N_Y \cdot T \cong N_Y \wr \text{Sym}(s)$ and by Lemma (4.6) $S_{*^s Y} = *^s S_Y$. Thus $N_{*^s Y}/S_{*^s Y} = N_Y \cdot T/(S_Y * S_Y * \dots * S_Y) \cong (N_Y/S_Y) \cdot T \cong M_Y \cdot T \cong M_Y \wr \text{Sym}(s)$. So $M_{*^s Y} \cong M_Y \wr \text{Sym}(s)$ and this wreath product can be written out as $(M_Y * M_Y * \dots * M_Y) \cdot \text{Sym}(s)$; keeping track of the isomorphisms we see that each factor M_Y acts on the corresponding factor Y of $*^s Y$ while the subgroup isomorphic to $\text{Sym}(s)$ still permutes these factors. Therefore the identification $M_{*^s Y} \cong M_Y \wr \text{Sym}(s)$ yields the isomorphism $*^s Y \cong Y^{ls}$. $\square$

Irreducibles and Control Of Closure 4.11. We have determined the structure of the permutation $kM_{*^s Y}$ -module $k *^s Y$ and of the permutation $kM_{\Delta^s Y}$ -module $k\Delta^s Y$ for irreducible Y . Now we turn to finding conditions under which $*^s Y$ or $\Delta^s Y$ is a closed element of $\mathcal{S}^q$. The following elementary lemma from group theory will be useful.

Lemma 4.12. *Let α be a finite set and let the subsets α_i partition α . For each i let H_i be a subgroup of $\text{Sym}(\alpha_i) \leq \text{Sym}(\alpha)$ that does not fix an element of α_i . Then $C_{\text{Sym}(\alpha)}(H_1 \times H_2 \times \dots \times H_t) = C_{\text{Sym}(\alpha_1)}(H_1) \times C_{\text{Sym}(\alpha_2)}(H_2) \times \dots \times C_{\text{Sym}(\alpha_t)}(H_t)$.*

Proof. Let $g \in \text{Sym}(\alpha)$ be any element that centralizes H_i . Then the action of $\langle g \rangle$ on α will admit as an invariant subset the set of elements of α not fixed by H_i ; by assumption this set is $\text{supp } H_i = \alpha_i$, so $g \in \text{Sym}(\alpha_i) \times \text{Sym}(\alpha \setminus \alpha_i)$. This holds for each i and so $g \in \text{Sym}(\alpha_1) \times \text{Sym}(\alpha_2) \times \dots \times \text{Sym}(\alpha_t)$, from which it follows $C_{\text{Sym}(\alpha)}(H_1 \times H_2 \times \dots \times H_t) \leq C_{\text{Sym}(\alpha_1)}(H_1) \times C_{\text{Sym}(\alpha_2)}(H_2) \times \dots \times C_{\text{Sym}(\alpha_t)}(H_t)$. The reverse inclusion is trivial. $\square$

If $X \in \mathcal{S}^q$ we define the *closure*, written $c(X)$ as the set $\text{Fix}(Q_X)$ for a choice Q_X of Sylow p -subgroup of $\text{Stab}(X)$. Closure is well defined up to equivalence.

Corollary 4.13. *Submultiplicativity of Closure. Assume $X_1, X_2, \dots, X_t$ belong to $\mathcal{S}^q$, and are exact. Then $c(X_1 * X_2 * \dots * X_t) \leq c(X_1) * c(X_2) * \dots * c(X_t)$.*

Proof. Set $X := X_1 * X_2 * \dots * X_t$, and let $Q_i := Q_{X_i}$, and define also $Q = Q_1 \times \dots \times Q_t$. Then the p -group Q is contained in S_X , so we can take $Q \leq Q_X$. We assume the X_i are exact, so we can apply (4.12) with $H_i = Q_i$, and get, taking $G = G_X$ and $G_i = G_{X_i}$, that

$$H := C_G(Q) = C_{G_1}(Q_1) \times \dots \times C_{G_t}(Q_t).$$

Let $\Xi_i = \Xi_{X_i}$; then we have $\Xi_1 * \dots * \Xi_t \subseteq \Xi_X$, and therefore

$$H \cap (\Xi_1 * \dots * \Xi_t) \subseteq H \cap \Xi_X.$$

We claim that equality holds. Namely taking $x \in \Xi_X$ that commutes with $Q = Q_1 \times \dots \times Q_t$, then each orbit of x is contained in X_i for some i and therefore $x \in \Xi_1 * \dots * \Xi_t$. So $H \cap \Xi_X$ is contained in $H \cap (\Xi_1 * \dots * \Xi_t)$. This last set is equal to $(C_{G_1}(Q_1) \cap \Xi_1) * \dots * (C_{G_t}(Q_t) \cap \Xi_t)$. Hence

$$\text{Fix}_{\Xi_1}(Q_1) * \dots * \text{Fix}_{\Xi_t}(Q_t) = C_G(Q) \cap \Xi_X \supseteq C_G(Q_X) \cap \Xi_X = \text{Fix}_{\Xi_X}(Q_X),$$

which is what we needed. $\square$

Corollary 4.14. *Let $X_1, X_2, \dots, X_t$ be exact closed members of $\mathcal{S}^q$. Then $X_1 * X_2 * \dots * X_t$ is exact and closed.*

Proof. Write $X = X_1 * X_2 * \dots * X_t$. We have $c(X_1 * X_2 * \dots * X_t) \leq c(X_1) * c(X_2) * \dots * c(X_t) = X_1 * X_2 * \dots * X_t$; thus we must have $c(X_1 * X_2 * \dots * X_t) = X_1 * X_2 * \dots * X_t$ and X is closed.

Let $\alpha_j = \text{supp} X_j$, writing $\alpha = \bigcup \alpha_j$ and $Q_j = Q_{X_j}$. Then by assumption Q_j is fixed point free on α_j . We are free to assume $Q_1 * Q_2 * \dots * Q_t \leq Q_X$ and so Q_X is fixed point free on α , so X is exact. $\square$

Lemma 4.15. *Let X be a closed exact element of $\mathcal{S}^q$. If i is a power of p then $\Delta^i X$ is closed.*

When $|X| > 1$, then $\Delta^i X$ is closed if and only if i is a power of p .

Proof. By (4.14) $*^i X$ is closed for all $i > 1$. Moreover by definition $\Delta^i X \subset *^i X$. Therefore we always have $c(\Delta^i X) \subset c(*^i X) = *^i X$. In other words, letting $\Xi = \Xi_{\Delta^i X}$, we know that

$$\text{Fix}_\Xi(Q_{\Delta^i X}) \subseteq (*^i X).$$

We must determine when the fixed point are only the diagonal. We take $Q_{\Delta^i X} = Q_X \wr Q$ where Q is a Sylow p -subgroup of $\text{Sym}(i)$.

(a) Assume that i is a power of p ; then Q is transitive on the support sets of the base group, and it follows that $\text{Fix}_{*^i X} = \Delta^i X$ and therefore the fixed points of $Q_{\Delta^i X}$ in $*^i X$ are as well just $\Delta^i X$.

(b) Assume $|X| > 1$. Suppose i is not a power of p ; then Q is not transitive on the support sets of the base group. In this case it has fixed points which belong to $\Delta^r X * \Delta^{i-r} X$ for some $1 \leq r \leq i$ but not to $\Delta^i X$. So $\Delta^i X$ is not closed. $\square$

As a partial converse to the above we have the following.

Lemma 4.16. *Let X be an irreducible closed projective element of $\mathcal{S}^q$. If $i \geq p$ is a power of p then $\Delta^i X$ is not closed.*

Proof. By assumption $X = \Xi_X$, and since the group $\langle \Xi_X \rangle$ is transitive on $\text{supp} X$ it follows that X is irreducible.

Let $S = S_{\Delta^i X}$ and $Q = Q_{\Delta^i X}$, and write $X_\Delta = \Delta^i X = \Delta X_1 \dots \Delta X_i$ where $\text{supp} X_j = \alpha_j$ for disjoint α_j and $\alpha = \bigcup \alpha_j$. We can now apply (4.5) and get

$$S = (S_{X_1} * \dots * S_{X_i}) \cdot T,$$

where $T \cong \text{Sym}(i)$. By (4.2) T is not transitive on α . Taking Sylow p -subgroups we get

$$Q = (Q_{X_1} * \dots * Q_{X_i}) \cdot P = P,$$

for a Sylow p -subgroup P of T , since $Q_X = 1$ by assumption. Then X_Δ is irreducible by (3.12) but not exact, since $P \leq T$ is not transitive; nor is X_Δ projective; so X_Δ cannot be closed by (3.16). $\square$

Remark 4.17. We have gathered enough information to give sufficient conditions that given an irreducible fixed point set $Y \in \mathcal{S}^q$ the element $*^i Y$ or the element $\Delta^i Y$ is also a fixed point set. These conditions are set out in the next theorem.

Theorem 4.18. *Recall the definition of $\mathcal{S}^q$ in (3.2). Let $Y \in \mathcal{S}^q$ be an exact (in the sense of (3.15)) irreducible fixed point set. Let $s < p$ be an integer and let t be a power of p . Then*

- (1) $*^s Y$ is a fixed point set.

(2) $\Delta^t Y$ is an irreducible fixed point set.

Proof. Recall that Y is closed, as it is a fixed point set. By (4.14) the element $*^s Y$ is closed. We want to show that in addition the permutation $kM_{*^s Y}$ -module $k*^s Y$ admits a projective summand.

Assume first that $|Y| > 1$. Then by (4.10) the pair $(M_{*^s Y}, *^s Y)$ is permutation isomorphic to the pair $(M_Y \wr \text{Sym}(s), Y^{\wr s})$. Thus the existence of the projective summand is a consequence of Lemma (1.10).


Now assume that $|Y| = 1$. Then $*^s Y = \Delta^s Y$ and by (4.9), $(M_{\Delta^s Y}, \Delta^s Y)$ is permutation isomorphic to (M_Y, Y) . Hence by assumption the permutation module $k\Delta^s Y$ is projective as a module for $M_{\Delta^s Y}$. So $*^s Y$ is also a fixed point set in this case.

That $\Delta^t Y$ is irreducible comes from Lemma (3.12). By Lemma (4.15) the element $\Delta^t Y$ is closed, and we want to show that the permutation $kM_{\Delta^t Y}$ -module $k\Delta^t Y$ admits a projective summand. By Lemma (4.9) the pair $(M_{\Delta^t Y}, \Delta^t Y)$ is permutation isomorphic to the pair (M_Y, Y) . The existence of the projective summand follows by assumption. $\square$

5. STRUCTURE OF PRODUCTS OF DISTINCT SETS.

We have examined the structure afforded by products such as $*^s Y$, where $Y \in \mathcal{S}^q$. Let $X, Y \in \mathcal{S}^q$ with possibly $X \neq Y$. In this section we consider the product $X * Y$ and the permutation $kM_{X * Y}$ -module it affords. We also introduce the notion of ‘coprime’.

Properties of $M_{X * Y}$ 5.1. In order to determine the permutation structure of $X * Y$ we may assume as usual that X and Y have disjoint support and that $X * Y = X \times Y$. Then the subgroups $N_X * N_Y$, $S_{X * Y}$ and $S_X * S_Y$ are all contained in $N_{X * Y}$. Thus we have the diagram of subgroups



We also have $S_X * S_Y = S_{X * Y} \cap (G_X * G_Y) = S_{X * Y} \cap (N_X * N_Y)$. The direct product $M_X \times M_Y$ therefore satisfies $M_X \times M_Y = \frac{N_X}{S_X} \times \frac{N_Y}{S_Y} \cong \frac{N_X \times N_Y}{S_X \times S_Y} = \frac{N_X * N_Y}{S_X * S_Y} = \frac{N_X * N_Y}{S_{X * Y} \cap (N_X * N_Y)} \cong \frac{(N_X * N_Y) \cdot S_{X * Y}}{S_{X * Y}} \leq \frac{N_{X * Y}}{S_{X * Y}} =: M_{X * Y}$. This means that the direct product $M_X \times M_Y$ can be naturally identified with a subgroup of $M_{X * Y}$.

Convention 5.2. Let X and Y be as in the previous section. The groups M_X and M_Y are factor groups of subgroups of $\text{Sym}(\mathbb{N})$ and as yet the expression $M_X * M_Y$ is not defined. Therefore we shall use $M_X * M_Y$ to denote the subgroup

of M_{X*Y} which is naturally identifiable with the direct product $M_X \times M_Y$. This subgroup is described in the previous section.

Coprime Sets 5.3. Let X and Y be two elements of $\mathcal{S}^q$. Then up to equivalence $X \subset \Xi_{qm}^q$ and $Y \subset \Xi_{qn}^q$ for some positive integers m and n . Now suppose that X and Y have disjoint support, so that we may write $X * Y = X \times Y$. One way to view $X * Y$ is that it is a subset of the $\text{Sym}(qm) * \text{Sym}(qn)$ -space $\Xi_{qm}^q * \Xi_{qn}^q$. The subgroup $N_X * N_Y$ acts naturally on the Cartesian product $X * Y$ and affords the permutation $k(N_X * N_Y)$ -module $kX \otimes kY$. A second point of view is that $X * Y$ is a subset of the $\text{Sym}_{q(m+n)}$ -space $\Xi_{q(m+n)}^q$. In this case $X * Y$ is a permutation N_{X*Y} -space and affords the permutation kN_{X*Y} -module $k(X * Y)$. There is at least as much structure in the second instance as in the first, and indeed we have an isomorphism of $kN_X * kN_Y$ -modules

$$\text{Res}_{N_X * N_Y}^{N_{X*Y}}(k(X * Y)) \cong kX \otimes kY.$$

Quotienting out the stabilizers of the various actions produces an isomorphism of $k(M_X * M_Y)$ -modules

$$\text{Res}_{M_X * M_Y}^{M_{X*Y}}(k(X * Y)) \cong kX \otimes kY.$$

The second point of view is the one we have to work with in order to show that $X * Y$ is a fixed point set. However, we will be very interested in those cases where the second view adds no more structure to the first, which leads to the following definition.

Definition. We say that X and Y are **coprime** if $N_{X*Y} = N_X * N_Y$, or equivalently if $N_{X*Y} \leq G_X * G_Y$.

Proposition 5.4. Let X and Y be coprime elements of $\mathcal{S}^q$. Then

- (1) we have $S_{X*Y} = S_X * S_Y$
- (2) we have $M_{X*Y} = M_X * M_Y$.

Proof. By hypothesis we have $N_{X*Y} \leq G_X * G_Y$ and moreover $N_{X*Y} = N_X * N_Y$.

(1) Since $S_X * S_Y = S_{X*Y} \cap (G_X * G_Y)$ it follows that $S_{X*Y} = S_X * S_Y$.

(2) This holds because $M_X * M_Y$ is a subgroup of M_{X*Y} and $M_{X*Y} := N_{X*Y} / S_{X*Y} = (N_X * N_Y) / (S_X * S_Y) \cong (N_X / S_X) \times (N_Y / S_Y) \cong M_X * M_Y$ (see (5.1)). $\square$

Structure of Products of Coprime Sets 5.5. Thus if X and Y are coprime we must have $M_{X*Y} = M_X * M_Y$. In this case the isomorphism of permutation $k(M_X * M_Y)$ -modules

$$\text{Res}_{M_X * M_Y}^{M_{X*Y}}(k(X * Y)) \cong kX \otimes kY.$$

is in fact an isomorphism of permutation kM_{X*Y} -modules

$$k(X * Y) \cong kX \otimes kY.$$

Lemma 5.6. Suppose Y and Z are two coprime elements of $\mathcal{S}^q$ which are fixed point sets. Assume $Y * Z$ is closed. Then $Y * Z$ is a fixed point set.

Proof. Recall that fixed point sets are closed. By assumption $Y * Z$ is closed, so it is sufficient to show that the permutation kM_{Y*Z} -module $k(Y * Z)$ has a projective summand. The fact that Y and Z are coprime implies that $M_{Y*Z} = M_Y \times M_Z$ and that $k(Y * Z)$ is isomorphic to $kY \otimes kZ$. Thus it suffices to show that the kM_Y -module kY and the kM_Z -module kZ each admit projective summands. This is known since Y and Z are fixed point sets. $\square$

Lemma 5.7. *Suppose Y and Z are two coprime exact elements of $\mathcal{S}^q$ which are fixed point sets. Then $Y * Z$ is a fixed point set.*

Proof. By Corollary (4.14) we know that $Y * Z$ is closed, and we may apply (5.6). $\square$

Proposition 5.8. *Let $X, Y \in \mathcal{S}^q$. Then X and Y are coprime if and only if they do not have an irreducible factor in common.*

Proof. We may assume that X and Y have disjoint supports so that $X * Y = X \times Y$.

Suppose first that X and Y share the irreducible factor Z . By hypothesis we have $X = Z_X \times W_X$ and $Y = Z_Y \times W_Y$ where Z_X and Z_Y are equivalent to Z . Let $\alpha = \text{supp} Z_X$ and $\beta = \text{supp} Z_Y$.

Recalling the definition made in Section (4.2) let $t := g_{\alpha \text{supp } Z_Y} \in \text{Sym}(\mathbb{N})$ be the unique order-preserving involution that maps α onto $\text{supp } Z_Y$ and satisfies $\text{supp } t = \alpha \cup \beta$.

We may assume that $Z_Y = Z_X^t$. Now $X^t = Z_Y \times W_X$ and $Y^t = Z_X \times W_Y$ so $(X \times Y)^t = X^t \times Y^t = X \times Y$. It follows that $t \in N_{X * Y}$. However, $\alpha \subset \text{supp } X$ while $\alpha^t = \beta \subset \text{supp } Y$ so it is clear that t does not leave $\text{supp } X$ or $\text{supp } Y$ invariant. Thus $t \notin G_X \times G_Y$, and X and Y are not coprime.

Suppose instead that X and Y are not coprime and let $t \in N_{X * Y}$ with $t \notin G_X * G_Y$. Let $X = X_1 \times X_2 \times \cdots \times X_s$ and $Y = Y_1 \times Y_2 \times \cdots \times Y_r$ be decompositions into irreducibles. Then $X \times Y = X_1 \times X_2 \times \cdots \times X_s \times Y_1 \times Y_2 \times \cdots \times Y_r$ is a decomposition into irreducibles. If for every $1 \leq i \leq s$ there is j such that $X_i^t = Y_j$ then t leaves $\text{supp } X$ and hence $\text{supp } Y$ invariant. By assumption this is not the case. Therefore by Corollary (3.10) for some $1 \leq i \leq s$ and $1 \leq j \leq r$ we have $X_i^t = Y_j$. Thus $X_i = Y_j$ modulo equivalence and X and Y share the irreducible factor X_i . $\square$

Theorem 5.9. *Suppose that $Y_1, Y_2, \dots, Y_t \in \mathcal{S}^q$ is a list of pairwise distinct irreducible exact fixed point sets and let $s_1, s_2, \dots, s_t$ be positive integers satisfying $s_i < p$ for every i . Then $Y_1^{s_1} * Y_2^{s_2} * \cdots * Y_t^{s_t}$ is a fixed point set.*

Proof. This follows directly from Theorem (4.18) and Lemma (5.7) because distinct irreducible elements of $\mathcal{F}$ are coprime. $\square$

To show how the assumption of exactness is necessary in the previous theorem we have the following.

Lemma 5.10. *Let Y and Z be closed projective elements of $\mathcal{S}^q$. Then $Y * Z$ is not closed.*

Proof. Write $X = Y * Z$. By assumption we have $Y = \Xi_Y$ and $Z = \Xi_Z$. In particular Y and Z are irreducible. If $Y \neq Z$ then Y and Z are coprime and we have $S_X = S_Y * S_Z$ by (5.4). Otherwise $Y = Z$ and again $S_X = S_Y * S_Z$ by (4.6). Accordingly we may always choose $Q_X = Q_Y * Q_Z$: but by assumption this gives $Q_X = 1$. So if X is closed then $X = \Xi_X$ which is a contradiction since Ξ_X is irreducible. $\square$

6. DECOMPOSING THE FIXED POINT SETS.

We have seen how to compose fixed point sets from irreducible ones, and in this section we will confirm that such compositions are the general form of fixed point sets. The results in this section will tend to be the converses of those in the previous section. By the end of the section the problem will have been reduced to finding the irreducible fixed point sets, which we will do in the next and final section.

Divisors of Fixed Point Sets 6.1. Given a fixed point set Z consider a decomposition of Z into irreducible factors - our goal is to show that each of the factors is again a fixed point set. This reduces the problem to determining all the irreducible fixed point sets.

Notation 6.2. If $X \in \mathcal{S}^q$ let $\tilde{Q}_X := \langle Q_X, X \rangle$ be the subgroup of G_X generated by Q_X and X . See Section (3.3) for the definitions of Q_X and X and Section (3.2) for the definition of $\mathcal{S}^q$.

Control Of Closure 6.3. In Section 3 we used Lemma (3.8) to split the task of showing that a set Y is a fixed point set into two parts. Thus to show that Y is a fixed point set we show that (i) Y is closed, and that (ii) the permutation kM_Y -module Y admits a projective summand. Since Lemma (1.7) implies that this second property passes readily to any factors in a decomposition of Y , the first results of this chapter deal with whether or not such factors are closed.

Remark 6.4. Suppose that Y and Z are elements of $\mathcal{S}$ such that $Y * Z$ is closed. We shall see in the proof of the next lemma that if $S_{Y*Z} = S_Y * S_Z$ then we may infer that Y and Z are closed. A sufficient condition for $S_{Y*Z} = S_Y * S_Z$ to hold is that the subgroups $\langle Y * Z \rangle$ and $\langle Y \rangle * \langle Z \rangle$ are equal. However, we shall take another approach and split this problem into two cases.

Lemma 6.5. *Suppose that Y and Z are coprime elements of $\mathcal{S}$ such that $Y * Z$ is closed. Then Y and Z are closed.*

Proof. The coprime condition gives $Q_{Y*Z} = Q_Y * Q_Z$. Thus $Y * Z = c(Y * Z) = \text{Fix}(Q_{Y*Z}) = \text{Fix}(Q_Y * Q_Z) \supset \text{Fix}(Q_Y) * \text{Fix}(Q_Z)$, giving $\text{Fix}(Q_Y) = Y$ and $\text{Fix}(Q_Z) = Z$. $\square$

Lemma 6.6. *Suppose Y is an irreducible element of $\mathcal{S}^q$ such that Y^i is closed. Then Y is closed.*

Proof. If $|Y| = 1$ then $Y^i = \Delta^i Y$. This case is dealt with in Lemma (7.10): no circularity of argument will result from this expedience. By Lemma (4.6) we have $S_{Y^i} = S_Y^i$, and so $Q_{Y^i} = Q_Y * Q_Y * \cdots * Q_Y$. The proof is now the same as the proof of the last lemma. $\square$

Corollary 6.7. *Suppose that Y and Z are coprime elements of $\mathcal{S}$ such that $Y * Z$ is a fixed point set. Then Y and Z are fixed point sets.*

Proof. The coprime condition gives $M_{Y*Z} = M_Y \times M_Z$ and $k(Y * Z) \cong kY \otimes kZ$, by Section (5.5). By hypothesis the permutation kM_{Y*Z} -module $k(Y * Z)$ admits a projective summand. Thus by Corollary (1.7) the permutation kM_Y -module kY and the permutation kM_Z -module kZ each admit projective summands. By Lemma (6.5) Y and Z are closed so Lemma (3.8) implies that Y and Z are fixed point sets. $\square$

Note that by Lemma (3.9) the decomposition in the statement of the following theorem is unique.

Theorem 6.8. *Let $Y \in \mathcal{S}^q$ be a fixed point set and write $Y = Y_1^{a_1} * Y_2^{a_2} * \cdots *$*

$Y_t^{a_t}$ where the Y_i are pairwise distinct irreducible elements and the a_i are positive integers. Then each Y_i is a fixed point set.

Proof. The list $Y_1^{a_1}, Y_2^{a_2}, \dots, Y_t^{a_t}$ is a list of pairwise coprime elements of $\mathcal{S}$ so Theorem (5.9) implies that each $Y_i^{a_i}$ is a fixed point set. In particular $Y_i^{a_i}$ is closed for every i .

Assume first $|Y_i| > 1$ for a given i . Now Lemma (4.10) tells us that for each i $kY_i^{a_i} \cong Y_i^{la_i}$ under the identification $kM_{Y_i^{a_i}} = kM_{Y_i} \wr \text{Sym}(a_i)$. Hence Lemma (1.9) asserts that the permutation kM_{Y_i} -module Y_i admits a projective summand. Now assume $|Y_i| = 1$; then $Y_i^{a_i} = \Delta^{a_i} Y_i$ and we can apply (4.9) which shows that the relevant permutation module has a projective summand. So in both cases, by (3.8) each Y_i is a fixed point set.

Meanwhile Lemma (6.5) and Lemma (6.6) tell us that each Y_i is closed, so by Lemma (3.8) Y_i is a fixed point set. $\square$

7. DETERMINING THE IRREDUCIBLE FIXED POINT SETS.

In this section we shall determine the structure of the irreducible fixed point sets (in the context of the present problem) and in doing so we will finally complete our description of a general fixed point set. The results of the previous sections will combine with the following propositions in order to arrive at the main theorem.

Reducing the Irreducible Fixed Point Sets 7.1. We will show in Corollary (7.6) that if Y is an irreducible fixed point set then $\hat{Q}$ must be transitive on $\text{supp } Y$. Thus $Q_Y := \langle Q_Y, Y \rangle$ transitively permutes the orbits on $\text{supp } Y$ of $\langle Y \rangle$. It will follow that $Y = \Delta^i X$ for some i and $X \in \mathcal{S}^q$, where the size of $\text{supp } X$ is equal to the size of a $\langle Y \rangle$ -orbit on $\text{supp } Y$. In Theorem (7.14) we find a bound for the orbit size, reducing the problem to the study of a finite, and in fact a very small, number of cases.

Lemma 7.2. *Suppose that $Y \in \mathcal{S}^q$ is closed irreducible. Suppose further that $Y = \Delta^s Z$ for some $Z \in \mathcal{S}^q$. Then Z is closed irreducible.*

Proof. The congruence $(X_1 * X_2) \Delta (X_1 * X_2) \cong (X_1 \Delta X_1) * (X_2 \Delta X_2)$ holds generally for X_1, X_2 in $\mathcal{S}^q$ so Z must be irreducible.

By Lemma (4.5) we have $S_Y = S_{\Delta^s Z} = S_Z \cdot T = {}^s S_Z \cdot T \cong S_Z \wr \text{Sym}(s)$ where $T \cong \text{Sym}(s)$ is the subgroup defined in (4.2) that permutes the factors of the base subgroup ${}^s S_Z \leq S_Y$. It follows from this that $Q_Y = {}^s Q_Z \cdot R$ where $R \leq T$ is a Sylow p -subgroup of T . Let $z \in c(Z)$ - then $c \in \text{Fix } Q_Z$. Clearly $y := z * z * \dots * z \in \text{Fix } {}^s Q_Z$; but y is also stabilized by T . Thus $y \in \text{Fix } {}^s Q_Z \cdot T \subset \text{Fix } Q_Y = Y$, and $z \in Z$. So Z is closed. $\square$

Orbits and Factorizations 7.3. If $G := \text{Sym}(m)$ is the Symmetric group on $m > 0$ letters and $H \leq G$ is a subgroup of G which leaves invariant the two subsets $\alpha \subset \{1, 2, \dots, m\}$ and $\beta \subset \{1, 2, \dots, m\}$, and if these subsets satisfy (i) $\alpha \cap \beta = \emptyset$ and (ii) $\alpha \cup \beta = \{1, 2, \dots, m\}$, then we know that $H \leq G$ sits inside the Young subgroup $\text{Sym}(\alpha) * \text{Sym}(\beta) \leq G = \text{Sym}(m)$. However, it is not necessarily true that there is a subgroup $H_\alpha \leq \text{Sym}(\alpha)$ of $\text{Sym}(\alpha) \leq G$ and a subgroup $H_\beta \leq \text{Sym}(\beta)$ of $\text{Sym}(\beta) \leq G$ for which $H = H_\alpha * H_\beta$.

The next theorem finds conditions which do allow us make this assumption.

Let $\tilde{Q}_X = \langle Q_X, X \rangle$.

Theorem 7.4. *Assume $X \in \mathcal{S}^q$ is exact and closed. Assume the group $\tilde{Q}_X$ has orbits $\lambda_1, \lambda_2, \dots, \lambda_s$ on the support of X . Then*

(1) $X = X_1 * X_2 * \dots * X_s$ where X_i has support λ_i and is closed as a subset of Ω_{X_i} and X_i exact.

(2) $Q_X = Q_1 * Q_2 * \dots * Q_s$ where $Q_i = Q_{X_i}$.

Proof. It suffices to consider $s = 2$. Write Q and $\tilde{Q}$ for Q_X and $\tilde{Q}_X$. We have $\tilde{Q} \subseteq H_1 \times H_2$ where $H_i := \text{Sym}(\lambda_i)$. Let π_i be the projection onto H_i , and set

$$Q_i := Q_{\pi_i}, X_i := X_{\pi_i}.$$

Then we have

(i) $Q \subseteq Q_1 \times Q_2$,

(ii) $X \subseteq X_1 * X_2$.

We claim that Q_i fixes each element of X_i . Namely, take $u_1 \in Q_1$, then choose $u_2 \in Q_2$ such that $u := u_1 u_2 \in Q$. Next, take $x_1 \in X_1$, and take $x_2 \in X_2$ such that $x := x_1 * x_2$ belongs to X . Then we have

$$x = x^u = (x_1^{u_1}) * (x_2^{u_2})$$

and $x_1 = x^{u_1}$. This shows that $Q_1 \subseteq S_{X_1}$, and similarly $Q_2 \subseteq S_{X_2}$.

Now take R_i to be a Sylow p -subgroup of S_{X_i} , containing Q_i . Then $R_1 \times R_2$ fixes each element of $X_1 * X_2$ and in particular it fixes each element of X . So it follows that the order of $R_1 \times R_2$ is $\leq$ the order of some Sylow p -subgroup of S_X and hence $\leq |Q|$.

On the other hand, from $Q \leq Q_1 \times Q_2 \leq R_1 \times R_2$ we have $|Q| \leq |R_1 \times R_2|$ and it follows that

$$Q = Q_1 \times Q_2 = R_1 \times R_2$$

and $Q_i = R_i$ which we can take for Q_{X_i} . This completes the proof of part (2).

It remains to show that $X_1 * X_2 \subseteq X$. Let $\Omega_i = \Xi_{X_i}$, then $\Omega_1 * \Omega_2 \subseteq \Xi_X$. We have

$$X_1 * X_2 \subseteq \text{Fix}_{\Omega_1}(Q_1) * \text{Fix}_{\Omega_2}(Q_2) \subseteq \text{Fix}_{\Omega_1 * \Omega_2}(Q) \subseteq \text{Fix}_{\Xi_X}(X);$$

and $\text{Fix}_{\Xi_X}(Q) = X$ since X is closed. So $X = X_1 * X_2$ and $X_i = \text{Fix}_{\Omega_i}(Q_i)$, that is X_i is closed. $\square$

Corollary 7.5. *Suppose that $X \in \mathcal{S}^q$ is closed, exact and irreducible. Then $\langle Q_X, X \rangle$ is a transitive subgroup of $G_X = \text{Sym}(\text{supp } X)$.*

Proof. This follows immediately from the previous Theorem (7.4). $\square$

Corollary 7.6. *Suppose that $X \in \mathcal{S}^q$ is closed, exact and irreducible. Let $\lambda_1, \lambda_2, \dots, \lambda_t$ be the orbits on $\text{supp } X$ of $\langle X \rangle$. Then Q_X transitively permutes the λ_i . In particular these orbits have the same size.*

Proof. Firstly the subgroup $\langle Q_X, X \rangle \leq G_X$ normalizes its subgroup $\langle X \rangle$ so $\langle Q_X, X \rangle$ permutes the λ_i , and it must do so transitively since $\langle Q_X, X \rangle$ is transitive on $\text{supp } X$ (by Corollary (7.5)). Now $\langle X \rangle$ acts trivially on the λ_i and it follows that Q_X must transitively permute the λ_i . $\square$

Transitive Sets 7.7. Let $X \in \mathcal{S}^q$. We say that X is a **transitive** set if $\langle X \rangle$ is transitive in its action on $\text{supp } X$.

Corollary 7.8. *Suppose that $X \in \mathcal{S}^q$ is closed, exact and irreducible. Then*

$X = \Delta^p Y$ for some integer $i \geq 0$ and some transitive set $Y \in \mathcal{S}^q$.

Proof. Let $\lambda_1, \lambda_2, \dots, \lambda_t$ be the orbits of $\langle X \rangle$ on $\text{supp } X$ - then the λ_i have equal sizes, by (7.6). Put $\lambda := \lambda_1$. Furthermore we have

$$X \leq X_{\lambda_1} * X_{\lambda_2} * \dots * X_{\lambda_t} = {}^* X_{\lambda}.$$

Now Q_X stabilizes X and transitively permutes the λ_i . It follows that $X \leq X_{\lambda_1} \Delta X_{\lambda_2} \Delta \dots \Delta X_{\lambda_t} = \Delta^t X_{\lambda}$, from which it follows that $X = \Delta^t Y$ where $Y := (X)\pi_{\lambda}$. Also, we see that t must be a p -power, for t is the size of the set $\{\lambda_1, \lambda_2, \dots, \lambda_t\}$, and the elements of this set are transitively permuted by Q_X . Now λ is an $\langle X \rangle$ -orbit on $\text{supp } X$ so $\langle (X)\pi_{\lambda} \rangle$ acts transitively on λ , whence Y is a transitive set. $\square$

Lemma 7.9. *Let $X \in \mathcal{S}^q$ be closed and exact. Then ZQ_X contains an element of order p fixed point free on $\text{supp } X$.*

Proof. By Lemma (7.4) it suffices to consider the case that $\tilde{Q}_X := \langle Q_X, X \rangle \leq G_n = \text{Sym}(2n)$ is transitive in its action on $\text{supp } X$. If $u \in ZQ_X$ is an element of ZQ_X then u centralizes Q_X and, since $u \in Q_X$, u centralizes $X = \text{Fix}(Q_X)$. Thus Q_X and X permute the fixed points on $\text{supp } X$ of $\langle u \rangle$ and in fact the transitive subgroup $\tilde{Q}_X$ permutes these fixed points. It follows that if we choose $u \in ZQ_X$ so that u has order p then u cannot fix any points in $\text{supp } X$. $\square$

7.10. Now let Y be an irreducible exact fixed point set in $\mathcal{S}^q$. By (7.9) there is some z central in Q_Y which is fixed point free and has order p .

We take a Sylow p -subgroup P of $C_{G_Y}(z)$ which contains Q_Y . Then P is even a Sylow p -subgroup of G_Y (as P is of the form $C_p \wr R$ for a Sylow subgroup of $\text{Sym}(s)$). Then by (2.2) there is some $y \in Y$ such that $Q_Y = \text{Stab}_P(y)$. We will use these elements z and y below.

Proposition 7.11. *Let Y be a closed element of $\mathcal{S}^q$. Suppose that Q_Y contains subgroups $R_1, R_2, \dots, R_l$. Let $\text{supp } R_i = \alpha_i$ and suppose that the α_i are disjoint. Suppose also that $*R_i \leq Q_Y$. Then $Y \subseteq *\text{Sym}(\alpha_i)$.*

Proof. Apply (4.12) with $H_i = R_i$. $\square$

Convention 7.12. Let $Y \in \mathcal{S}^q$ be an exact fixed point set. By Lemma (7.9) there is an element $z \in ZQ_Y$ which has order p and which is fixed point free on $\text{supp } Y$. Taking the cycle decomposition of z we have $z = z_1 * z_2 * \dots * z_s$ where each z_i is a p -cycle. Let P_z be a Sylow p -subgroup of $C_{G_Y}(z)$. Now $C_{G_Y}(z)$ certainly centralizes z and must permute the $\{z_i\}$. Thus $B_z := \langle z_1 \rangle * \langle z_2 \rangle * \dots * \langle z_s \rangle$ is a normal p -subgroup of $C_{G_Y}(z)$ - in fact B_z is the base subgroup of the wreath product $C_{G_Y}(z) \cong C_p \wr \text{Sym}(s)$, as we have seen in (4.1). It follows that $B_z \leq P_z$.

By (4.1) we know that P_z is a Sylow p -subgroup of G_Y . We observe that in this context the roles of p and q have interchanged.

The next theorem determines the possible transitive irreducible exact fixed point sets. Recall that $d(Y) = |\text{supp } (Y)|$.

Theorem 7.13. *Assume that q is prime. Let $Y \in \mathcal{S}^q$ be an irreducible fixed point set which is exact and transitive. Let $d(Y) = |\text{supp}(Y)|$. Then one of the following holds*

(1) $d(Y) = pq$

(2) $d(Y) = q$.

In case (2), Y is the set of q -cycles in $G_Y \cong \text{Sym}(q)$.

Proof. (a) Let $z \in Z(Q_Y)$ be an element of order p which is fixed point free on $\text{supp}(Y)$. [This exists by (7.10).] Then $Q_Y \subseteq C_G(z)$. Take a Sylow p -subgroup $P := P_Z$ of $C_G(z)$ containing Q_Y . Then $P \in \text{Syl}_p(G)$. [To see this: we know $C_G(z) \cong C_p \wr \text{Sym}(s)$ where z has s p -cycles, and z has support equal to the total support of G_Y . So P is of the form $C_p \wr R$ where R is a Sylow subgroup of $\text{Sym}(s)$. On the other hand, a Sylow subgroup of G_Y is exactly of this form, since p divides the size of its support.]

Now we use (2.2), which gives us that there is some $y \in Y$ such that $Q_Y = \text{Stab}_P(y)$. Then z and y commute.

(b) Since y and z commute, conjugation by y permutes the cycles of z . Let I be an index set labelling the orbits, and write $\bar{z}_i$ for the product of the cycles in orbit i . Then $\bar{z}_i$ commutes with y , for each i . Recall from (7.12) that P contains each cycle of z , so $\bar{z}_i$ belongs to P and therefore it belongs to $\text{Stab}_P(y) = Q_Y$. Now we apply (7.11) with $R_i = \langle \bar{z}_i \rangle$, and we get

$$Y \subseteq \prod_i \text{Sym}(\text{supp}(\bar{z}_i)).$$

By the hypothesis, Y is transitive, and therefore I has size 1. Also $z = \bar{z}_i$. This has the following consequences.

(i) The orbit size must divide the order of $\langle y \rangle$, that is s divides q .

(ii) The group $\langle y, z \rangle$ is transitive on the support of Y .

Since z and y commute, conjugation by z also permutes the cycles of y .

(iii) By (ii) z must be transitive on the cycles of y and then we have that the number of cycles m of y divides the order of $\langle z \rangle$ which is p . But p is prime, so either $m = 1$ or $m = p$.

CASE 1 $m = p$. Then we have $\text{d1}(Y) = pq$.

CASE 2 $m = 1$. Then $\text{d1}(Y) = q = sp$. In this case, Y is a set of q -cycles in $G_Y \cong \text{Sym}(q)$. $\square$

Transitive Fixed Point Sets For Prime q 7.14. We assume in this section that q is a prime. Theorem (7.13) places a severe restriction on the possible degrees of the transitive fixed point sets. We shall now compute the fixed point sets with these degrees.

Firstly we consider the case p, q coprime. Thus Theorem (7.13) asserts that any irreducible exact transitive fixed point set has degree qp , which means we are dealing with the action of $\text{Sym}(qp)$ on its conjugacy class Ξ_{qp}^q of fixed point free elements of order q . Let $X \subset \Xi_{qp}^q$ be an irreducible exact transitive fixed point set and let $x \in X$. Then a corresponding vertex Q_X fixes x and so $Q_X \leq C_{\text{Sym}(qp)}(x) \cong \mathbb{Z}_q \wr \text{Sym}(p)$. It follows that the size of Q_X is 1 or p , whence it must be p since X is transitive. Thus Q_X is generated by an element of order p which we denote by z in order to be consistent with the statement of Corollary (7.10). If x is written as the p -fold product of q -cycles

$$x = (a_{11}, a_{12}, \dots, a_{1q})(a_{21}, a_{22}, \dots, a_{2q}) \dots (a_{p1}, a_{p2}, \dots, a_{pq})$$

then z may be taken to be the q -fold product of p -cycles

$$z = (a_{11}, a_{21}, \dots, a_{p1})(a_{12}, a_{22}, \dots, a_{p2}) \dots (a_{1q}, a_{2q}, \dots, a_{pq}),$$

and $X = C_{\text{Sym}(qp)}(Q_X) \cap \Xi_{qp}^q = C_{\text{Sym}(qp)}(z) \cap \Xi_{qp}^q$.

Now we consider the case $p = q$. In this case Theorem (7.13) asserts that any transitive fixed point set has degree p or p^2 . To find a transitive fixed point set X with $d(X) = p$ we must consider the action of $\text{Sym}(p)$ on its conjugacy class Ξ_p^p of fixed point free elements of order p . Now a Sylow p -subgroup of $\text{Sym}(p)$ has order p so a corresponding vertex Q_X must have order p , since $Q_X = 1$ is ruled out by the transitivity of X . Furthermore $X = C_{\text{Sym}(qp)}(Q_X) \cap \Xi_p^p = Q_X \cap \Xi_p^p = Q_X \setminus 1$.

To find a transitive fixed point set X with $d(X) = p^2$ we must consider the action of $\text{Sym}(p^2)$ on its conjugacy class $\Xi_{p^2}^p$ of fixed point free elements of order p . Take $x \in \Xi_{p^2}^p$ and let $C_x := C_{\text{Sym}(p^2)}(x) \cong \mathbb{Z}_p \wr \text{Sym}(p)$. Now C_x contains a Sylow p -subgroup P of $\text{Sym}(p^2)$ and we may assume that X is such that $Q_X \leq P \leq C_x$, so $x \in X$. By Corollary (7.10) there is $y \in X$ such that $Q_X = \text{Stab}_P(y)$; hence Q_X fixes both x and y , and also $x \in Q_X$ since $x \in \text{Stab}_P(y)$. In the proof of Theorem (7.13) we used the fact that since y commutes with x it must induce a permutation of the p p -cycles that compose x , and by a remark in the third paragraph of that proof we see that this induced action must be transitive, a condition on y that we will refer to as condition (T). It can be shown that for all y satisfying condition (T) we have $\text{Stab}_{C_x}(y) = \langle x, y \rangle$ and so $Q_X \leq \langle x, y \rangle$. Thus we either have (i) $Q_X = \langle x \rangle$ or (ii) $Q_X = \langle x, y \rangle$. With (ii) it at first appears that different y might yield non-conjugate subgroups $\langle x, y \rangle$, but in fact it may be shown that any two elements y_1 and y_2 in $\Xi_{p^2}^p$ satisfying condition (T) are conjugate in C_x , and the resulting subgroups $\langle x, y_1 \rangle$ and $\langle x, y_2 \rangle$ will be conjugate in C_x . The corresponding fixed point sets are easily computed.

We see that when $p \neq q$ there is essentially at most one transitive fixed point set, and when $p = q$ there are at most three essentially different fixed point sets. In either case the number of transitive fixed point sets is certainly finite.

Definition 7.15. Assume that H is any finite group and Ω is any H -set. Assume that the permutation module $k\Omega$ has a projective summand.

Define $\kappa(\Omega)$ to be the lowest positive integer u such that the permutation module $(k\Omega)^{lu}$ for $H \wr \text{Sym}(u)$ does not have a projective summand, allowing the possibility $\kappa(\Omega) = \infty$ if no such u exists. Recall (1.10) from which we get $\kappa(\Omega) \geq p$. If $j > \kappa(\Omega)$ then $k\Omega$ does not admit a projective summand by (1.11).

Example 7.16. Let $H = C_2$, cyclic of order 2, and let $p = 3$.

(a) Take $M = k\Omega$ with $\Omega = H$ where H acts by left multiplication. Then consider M^{i3} , a module for $H \wr \text{Sym}(3)$.

If $\langle \tau \rangle = H$ then take the basis $\{1, \tau\}$ of M , and take the basis for $M^{\otimes 3}$ consisting of all $v_1 \otimes v_2 \otimes v_3$ with $v_i = 1$ or τ . Take a 3-cycle g of $\text{Sym}(3)$. This has two orbits of size 3 on this basis and two orbits of size 1. It follows that M^{i3} must have a projective summand as a module for $\langle g \rangle$. But $\langle g \rangle$ is a Sylow p -subgroup of $G := H \wr \text{Sym}(3)$ and we deduce that M^{i3} must have a projective summand as a module for G .

So $\kappa(\Omega) = 2 = p - 1$.

(b) Now take $M = k$, the trivial module, so that Ω is a set with one element on which H acts trivially. Then M^{ia} is 1-dimensional for all $a \geq 1$ and hence does not have a projective summand for G . So $\kappa(\Omega) = \infty$.

Definition 7.17. We shall call a set $X \in \mathcal{S}^q$ **projective-free** if none of its

irreducible components is projective. By (3.15) products of exact sets are exact so projective-free implies exact.

Recall from (2.13) that if $Z \in \mathcal{S}^q$ is closed then Z is a fixed point set if and only if the kM_Z -module kZ admits a projective summand.

Theorem 7.18. *Assume the setup of before, with elements of $\mathcal{S}^q$. Assume $X \in \mathcal{S}^q$.*

- (1) *X is an irreducible exact fixed point set $\Leftrightarrow X = \Delta^{p^i} Y$ where Y is a transitive irreducible exact fixed point set and $i \geq 0$.*
- (2) *X is a projective-free fixed point set $\Leftrightarrow X = X_1^{a_1} * \cdots * X_t^{a_t}$ where the X_i are pairwise coprime irreducible exact fixed point set, and $1 \leq a_i < \kappa(X_i)$.*
- (3) *X is a fixed point set $\Leftrightarrow X = W * V$ where W is a projective-free fixed point set and V is an irreducible projective fixed point set.*

Proof. (1) ‘ $\Rightarrow$ ’ By (7.8) $X = \Delta^{p^i} Y$ for Y transitive (and therefore irreducible). From (7.2) Y is closed, so by (4.16) it cannot be projective; and by (3.16) it must be exact. Also kY admits a projective summand by (4.9): so Y is a fixed point set.

‘ $\Leftarrow$ ’ By (4.18) X is an irreducible fixed point set. The case $i = 0$ is trivial so assume $i > 0$. By (4.5) it is clear that $Q_X \neq 1$, so X is not projective and must be irreducible.

(2) ‘ $\Rightarrow$ ’ Let $X = X_1^{a_1} * \cdots * X_t^{a_t}$ be a decomposition into irreducibles. By (6.8) each X_j is a fixed point set. But $X_j^{a_j}$ is also a fixed point set by (6.7) applied to X so by (4.10) $1 \leq a_j < \kappa(X_j)$. By assumption each X_j is not projective, hence exact.

‘ $\Leftarrow$ ’ We have X and each $X_j^{a_j}$ is exact and closed by (4.14). By assumption $1 \leq a_j < \kappa(X_j)$ so (4.10) tells us that each $X_j^{a_j}$ is a fixed point set. It follows by (5.7) that X is a fixed point set.

(3) ‘ $\Rightarrow$ ’ Factorize X into a product of irreducible sets. Say

$$X = W_1^{a_1} * \cdots * W_s^{a_s} * V_1^{b_1} * \cdots * V_s^{b_s},$$

and write $W = W_1^{a_1} * \cdots * W_s^{a_s}$ and $V = V_1^{b_1} * \cdots * V_s^{b_s}$. Assume the W_i are exact and the V_j are projective. If there is no projective factor then there is nothing to do. So assume that $t \geq 1$. By (6.7) V and W are fixed point sets. We wish to show that $t = 1$ and $b = 1$. By (6.7) each $V_j^{b_j}$ is a fixed point set and must be closed, which by (5.10) means that $b_j = 1$. Now V is a fixed point set, hence closed, so by (5.10) we get $t = 1$.

‘ $\Leftarrow$ ’ We first show that X is closed. Since V is projective, we know that $V = \Xi_V$ and because W and V are necessarily coprime we know by (5.4) that $Q_X = Q_W * Q_V = Q_W * 1$. The product W is projective-free, hence exact, which means that Q_W is fixed point free on $\text{supp} Q_W$. We can now compute directly that

$$\text{Fix}_{\Xi_{W*V}}(Q_W * 1) = W * V,$$

and X is closed. It follows from (5.6) that X is a fixed point set. $\square$

REFERENCES

- [1] M. Broué On Scott modules and p -permutation modules: an approach through the Brauer morphism. *Proc. Amer. Math. Soc.* 93 (1985), no. 3, 401–408.
- [2] P. Collings Fixed point free involutions over a field of characteristic two and other actions of the symmetric group by conjugation on its own elements. *PhD Thesis* Oxford University 2004
- [3] J. A. Green On the indecomposable representations of a finite group. *Math. Z.* 70 1958/59 430–445.

- [4] N. F. J. Inglis, R. W. Richardson and J. Saxl An explicit model for the complex representations of S_n . *Arch. Math. (Basel)* 54 (1990), no. 3, 258–259.
 - [5] G. James and A. Kerber The representation theory of the Symmetric group. *Encyclopedia of Mathematics and its Applications*, 16 Addison-Wesley Publishing Co., Reading, Mass., 1981.
 - [6] J. Thévenaz. G -algebras and modular representation theory. *Oxford Mathematical Monographs*, Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1995.
-